

NOTES ON EXISTENCE OF CLASSICAL SOLUTIONS OF SECOND ORDER ELLIPTIC EQUATIONS

DANIEL BALLESTEROS-CHÁVEZ

May 10, 2025

CONTENTS

1. Introduction	1
2. Ellipticity	3
3. Maximum principle for the Laplace equation	5
4. The Poisson's equation in domains of \mathbb{R}^n	6
4.1. Perron's Method	9
4.2. Interior estimates	11
4.3. Regularity of solutions to the boundary value problem: Kellogg's Theorem	11
4.4. Harnack's Inequality	12
5. Linear Second Order Partial Differential Equations: Schauder Theory.	12
6. Quasi-linear Second Order Partial Differential Equations: The Leray-Schauder Fixed Point Theorem.	17
7. Fully Nonlinear Partial Differential Equation	18
8. Maximum Principle for Nonlinear PDE's	19
8.1. The classical Aleksandrov-Bakelman-Pucci Theorem	20
8.2. Krylov-Safonov theorem	21
8.3. Evans - Krylov	21
References	21

1. INTRODUCTION

Be aware that this is still work in progress. Any comment is more than welcome.

These notes are based strongly in the books of D. Gilbarg and N. S. Trudinger [3], L. Nirenberg [5], L. C. Evans [2]. The aim is to give a quick introduction to the a priori estimates approach to prove existence of solutions for elliptic second order equations.

When we start the study of elliptic equations, the main examples are the Laplace and Poisson's equations. For these one can establish the maximum principle and use it to estimate on the values of the solutions and of their higher order derivatives. For more general equations, a maximum principle can also be established if some conditions on the coefficients of the differential operator are satisfied. Then, even without explicit expression for their solutions, it is possible to get similar inequalities for their values and get some regularity results.

The main idea is that, using topological theory on Banach spaces and functional analysis it is possible to get existence results for solutions of the Dirichlet problem based on these inequalities. In the case of linear equations, the Schauder estimates implies the existence results for a wide variety of problems depending on the properties of the coefficients of the differential operator. Then by means of the continuity method, one can establish the invertibility of such operator from a known one.

For quasilinear equations, the problem is more complicated, and topological methods like degree theory and fixed point results are extended to infinite dimensional Banach spaces. The Leray-Schauder fixed point theorem in Banach spaces gives the existence theory for the Dirichlet problem of a large class of quasilinear equations.

Finally we say something about the fully nonlinear case and how continuity method can be also applied if certain a priori estimates are established.

These notes are not exhaustive, and there are many different results regarding the influence of the geometry of the domain Ω and its boundary $\partial\Omega$ to the solution of the Dirichlet problem that are not mentioned.

We will study second order partial differential equations that can be linear, or not, depending on some properties of its coefficients. For instance let $\Omega \subset \mathbb{R}^n$ be a domain (open and connected), $\mathbb{R}^{n \times n}$ the set of all square symmetric matrices n by n . A second order equation can be written in general as a function

$$F : \Omega \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$$

$$(1) \quad F[u] = F(x, u, Du, D^2u) = 0.$$

In case that the coefficients of highest order depend only on $(x, u, Du) \in \Omega \times \mathbb{R} \times \mathbb{R}^n$, then the equation is called **quasi-linear**, and can be written as

$$(2) \quad Qu = a^{ij}(x, u, Du)D_{ij}u + b(x, u, Du).$$

A quasi-linear equation such that the highest order term is linear is called **semi-linear**:

$$(3) \quad Pu = a^{ij}(x)D_{ij}u + b(x, u, Du).$$

If all coefficients are depending only on $x \in \Omega$ we refer to it as a **linear** equation

$$(4) \quad Lu = a^{ij}(x)D_{ij}u + b^i(x)D_iu + c(x)u.$$

A **fully nonlinear** equation is a second order equation F that cannot be written as a quasi-linear equation.

Quasilinear equations are called **elliptic** if the coefficient matrix $[a^{ij}]$ is positive definite, moreover, if λ and Λ denote the minimum and maximum eigenvalues of $[a^{ij}]$, then the equation is **elliptic** if $\lambda \geq 0$, **strictly elliptic** when $\lambda \geq \epsilon > 0$, and **uniformly elliptic** when the ratio Λ/λ is bounded. A fully nonlinear equation F is elliptic if the matrix $F_{ij} = \partial F / \partial u_{ij}$ is positive definite.

The **Hölder space** $C^{k,\alpha}(\bar{\Omega})$, with exponent $0 < \alpha < 1$ is the Banach space of functions f with norm

$$(5) \quad |f|_{C^{k,\alpha}(\Omega)} = |f|_{C^k(\Omega)} + \max_{|j|=k} |D^j f|_{C^\alpha(\Omega)},$$

where

$$(6) \quad |f|_{C^k(\Omega)} = \sum_{j=0}^k \max_{|r|=j} \sup_{x \in \Omega} |D^r f| \quad ; \quad |f|_{C^\alpha(\Omega)} = \sup_{\substack{x \neq y, \\ x, y \in \Omega}} \frac{|f(x) - f(y)|}{|x - y|^\alpha}.$$

Hölder spaces are Banach spaces with the norm (5) when functions are defined in an open bounded set Ω . Let $\Omega \subset \mathbb{R}^n$ a bounded domain, then the Hölder space $C^{k,\alpha}(\bar{\Omega})$ is a non-reflexive and non-separable Banach space. If $k_1 + \alpha_1 > k_2 + \alpha_2$ then the inclusion $C^{k_2,\alpha_2}(\bar{\Omega}) \subset C^{k_1,\alpha_1}(\bar{\Omega})$

The following semi-norms are also used:

Non-dimensional norms: If Ω is bounded and $d = \text{diam } \Omega$ we define:

$$(7) \quad |f|'_{C^k(\Omega)} = \sum_{j=0}^k d^j \max_{|r|=j} \sup_{x \in \Omega} |D^r f| \quad ; \quad |f|'_{C^{k,\alpha}(\Omega)} = |f|'_{C^k(\Omega)} + d^{k+\alpha} \max_{|j|=k} |D^j f|_{C^\alpha(\Omega)}.$$

Interior norms: $d_x = \text{dist}(x, \partial\Omega)$ and $d_{x,y} = \min\{d_x, d_y\}$:

$$(8) \quad |f|_{C^k(\Omega)}^* = \sum_{j=0}^k \max_{|r|=j} \sup_{x \in \Omega} d_x^j |D^r f| \quad ; \quad [f]_{C^{k,\alpha}(\Omega)}^* = \sup_{x \neq y; |x-y|=d_{x,y}} d_{x,y}^{k+\alpha} \frac{|D^k f(x) - D^k f(y)|}{|x - y|^\alpha}.$$

$$|f|_{C^{k,\alpha}(\Omega)}^* = |f|_{C^k(\Omega)}^* + [f]_{C^{k,\alpha}(\Omega)}^*.$$

Uniform Limit Theorem. Let X be a topological space and E a metric space. Consider the family $\{f_k\}_{k \in \mathbb{N}}$ of functions $f_k : X \rightarrow E$ and suppose it converges uniformly to a function $f : X \rightarrow E$. Then the following holds:

- If each f_k is continuous then the limit function f is also continuous.
- If each f_k is uniformly continuous then the limit function f is also uniformly continuous.

Arzela-Ascoli Theorem. This is a fundamental result in mathematical analysis. It gives the necessary and sufficient conditions for a sequence of functions to have a uniformly convergent subsequence.

Theorem 1.1 ([1], appx. D). *Suppose that $\{f_k\}_{k=1}^\infty$ is a sequence of functions defined on a domain $\Omega \subset \mathbb{R}^n$ such that*

- (1) *It is uniformly bounded, i.e. there is a constant $M > 0$ such that $|f_k(x)| \leq M$ for all $x \in \Omega$ and all k .*
- (2) *It is uniformly equicontinuous: for each $\epsilon > 0$, there exists $\delta > 0$ such that if $|x - y| < \delta$ then $|f_k(x) - f_k(y)| < \epsilon$, for $x, y \in \Omega$ and all k .*

Then there exists a sub-sequence $\{f_{k_j}\}_{j=1}^\infty \subset \{f_k\}_{k=1}^\infty$ and a continuous function f such that $f_{k_j} \rightarrow f$, uniformly on compact subsets of Ω .

Some of the applications of the Arzela-Ascoli theorem we will be using are presented in the following examples.

Example 1.2. Let $\{f_k\}$ be a uniformly bounded sequence of differentiable functions defined in a compact set $K \subset \mathbb{R}^n$. If $\{Df_k\}$ is uniformly bounded by $M > 0$, then by the mean value theorem we have

$$(9) \quad |f_k(x) - f_k(y)| \leq \sup_{z \in K} |Df_k(z)| |x - y| \leq M |x - y|.$$

Note that a very similar argument works if f_k is a sequence of Lipschitz functions with the same Lipschitz constant. This is equivalent to equicontinuity of the family. Then by the Arzela-Ascoli theorem the sequence f_k has a subsequence that converges uniformly on compact sets. The limit function is also Lipschitz with same constant M .

Example 1.3. Let $\{f_k\}$ be a uniformly bounded sequence of continuous functions defined in a compact set $K \subset \mathbb{R}^n$ and such that $f_k \in C^\alpha(K)$ for all k , and even more, there is a positive constant $M > 0$ such that $|f_k|_{C^\alpha(K)} \leq M$, for all k . Then again, this is equivalent to equicontinuity of the family and by the Arzela-Ascoli theorem the sequence f_k has a subsequence that converges uniformly on compact sets.

Even more, if $f_k \in C^{j,\alpha}(K)$, then the limit of the subsequence f_{k_i} is also in $C^{j,\alpha}(K)$ since

$$(10) \quad |D^j f(x) - D^j f(y)| \leq \lim_{i \rightarrow \infty} \max_{|r|=j} |D^r f_{k_i}(x) - D^r f_{k_i}(y)| \leq C|x - y|^\alpha.$$

Remark Note that in general if $\{f_k\}_{k \in \mathbb{N}}$ is a sequence of Lipschitz functions of constant M_n that converges uniformly to a function f , then f is not necessarily Lipschitz. Consider the example $f_k(x) = \sqrt{x + \frac{1}{k}}$ in the interval $[0, 1]$.

Theorem 1.4 ([1], T1 S9.2.1). Let X be a Banach space. Assume $A : X \rightarrow X$ is a mapping such that

$$(11) \quad \|A(x) - A(y)\| \leq c \|x - y\|,$$

for all $x, y \in X$ and for some constant $c < 1$. Then A has a unique fixed point.

Proof. For any $x_0 \in X$ we consider the following iterative scheme: $x_{k+1} := A(x_k)$. Then it follows that for $k = 1, 2, \dots$, we have

$$(12) \quad \|A(x_{k+1}) - A(x_k)\| \leq c^k \|A(x_0) - x_0\|.$$

Then, if $k \geq l$,

$$(13) \quad \|x_k - x_l\| \leq \|A(x_{k-1}) - A(x_{l-1})\| \leq \sum_{i=l-1}^{k-2} \|A(x_{i+1}) - A(x_i)\| \leq \|A(x_0) - x_0\| \sum_{i=l-1}^{k-2} c^i,$$

this shows that $\{x_k\}_{k=1}^\infty$ is a Cauchy sequence in X and hence there the limit $\lim_k x_k = x$, exists. Now that x is a fixed point of A follows from the iteration scheme and the continuity of A :

$$(14) \quad A(x) = \lim_{k \rightarrow \infty} A(x_k) = \lim_{k \rightarrow \infty} x_{k+1} = x.$$

The uniqueness follows from the assumption

$$(15) \quad \|A(x) - A(y)\| \leq c \|x - y\|.$$

□

The following implicit function theorem is very useful when studying the linearised operator of a nonlinear PDE. It is a way to prove there are solutions near a given one.

Theorem 1.5. Let B_1, B_2, X Banach spaces and let $\mathcal{G} : B_1 \times X \rightarrow B_2$. Let $(u_0, s_0) \in B_1 \times X$ such that

- $\mathcal{G}(u_0, s_0) = 0$,
- \mathcal{G} is C^1 near (u_0, s_0) .
- The Fréchet derivative of \mathcal{G} at (u_0, s_0) is invertible (a linear isomorphism between B_1 and B_2).

Then there exists a neighbourhood N of s_0 in X such that the equation $\mathcal{G}(u, s) = 0$, is solvable for each $s \in N$, with solution $u = u_s \in B_1$.

If $F : X \rightarrow B_1$ is defined as $s \mapsto u_s$ for $s \in N$, such that $\mathcal{G}(F(s), s) = 0$, then F is differentiable at s_0 . In particular there is a constant C such that

$$(16) \quad |F(s) - F(s_0)|_{B_1} \leq C|s - s_0|_X.$$

2. ELLIPTICITY

Consider the following example of partial differential operator in \mathbb{R}^2 :

$$F = a_{11}u_{11} + a_{12}u_{12} + a_{22}u_{22} + b_1u_1 + b_2u_2 + cu,$$

then, the matrix F^{ij} is given by

$$(F^{ij}) = \begin{pmatrix} a_{11} & \frac{a_{12}}{2} \\ \frac{a_{12}}{2} & a_{22} \end{pmatrix},$$

hence, F is elliptic if

$$\det(F^{ij}) = a_{11}a_{22} - \frac{a_{12}^2}{4} \geq 0,$$

or equivalently

$$4a_{11}a_{22} - a_{12}^2 \geq 0.$$

It is not a trivial matter to generalise the concept of ellipticity to a higher order system of N equations for N functions u_1, \dots, u_N . Here we present the notion of non-characteristics and how it can help us to understand ellipticity from this point of view.

In general one can consider the system of equations

$$F_i \left(x, u_1, \dots, u_N, \frac{\partial^k u_j}{\partial x_1^{k_1} \dots \partial x_n^{k_n}}, \dots \right) = 0,$$

with $i, j = 1, 2, \dots, N$, and $k = k_1 + \dots + k_n$ where $1 \leq k \leq n_j$, that is, for each function u_j there is a highest order derivative n_j appearing in the system.

The semilinear case. Consider the case where each F_i is linear in the highest order derivative, then we can write the i -th equation as

$$(17) \quad \sum_{j=1}^N \sum_{k_1+\dots+k_n=n_j} a_{ij}^{k_1 \dots k_n} \frac{\partial^{n_j} u_j}{\partial x_1^{k_1} \dots \partial x_n^{k_n}} + G_i \left(x, u_1, \dots, u_N, \frac{\partial^m u_j}{\partial x_1^{m_1} \dots \partial x_n^{m_n}}, \dots \right) = 0,$$

here, $1 \leq m \leq n_j - 1$.

Definition 2.1. Let $p \in \mathbb{R}^n$, and S be a smooth hypersurface containing p . Suppose that u_1, \dots, u_N are functions satisfying the system of semilinear system of equations (17). Suppose that the values of the functions u_j and the values of their derivatives up to order $n_j - 1$ are known on the surface in a neighbourhood around p . If we can calculate the n_j -th order derivatives of u_j at the point p , then the surface S is called **non-characteristic** at p with respect to the semilinear system.

Suppose that y_1, \dots, y_n are new coordinates around the point p , such that, around a neighbourhood of p , the surface S is given by $y_1 = 0$.

Observe that, since for the function u_j we know its $n_j - 1$ derivatives, then it is possible to obtain the n_j -th derivative in directions which are tangent to the surface.

Then the problem is now to compute the n_j -th order derivatives $\frac{\partial^{n_j} u_j}{\partial y_1^{n_j}}$. By the chain rule one sees that

$$(18) \quad \frac{\partial^{n_j} u_j}{\partial x_1^{k_1} \dots \partial x_n^{k_n}} = \frac{\partial^{n_j} u_j}{\partial y_1^{n_j}} \left(\frac{\partial y_1}{\partial x_1} \right)^{k_1} \dots \left(\frac{\partial y_1}{\partial x_n} \right)^{k_n} + \begin{array}{c} \text{derivatives of order } n_j \\ \text{mixed with} \\ \text{tangent directions} \end{array} + \text{lower order terms.}$$

Making use of the equation (17), we obtain

$$(19) \quad \sum_{j=1}^N \sum_{k_1+\dots+k_n=n_j} a_{ij}^{k_1 \dots k_n} \frac{\partial^{n_j} u_j}{\partial y_1^{n_j}} \left(\frac{\partial y_1}{\partial x_1} \right)^{k_1} \dots \left(\frac{\partial y_1}{\partial x_n} \right)^{k_n} + \dots + G_i = 0.$$

These equations can be solved for $\frac{\partial^{n_j} u_j}{\partial y_1^{n_j}}$ if and only if the determinant of the $N \times N$ matrix

$$(20) \quad m_{ij} = \sum_{k_1+\dots+k_n=n_j} a_{ij}^{k_1 \dots k_n} \left(\frac{\partial y_1}{\partial x_1} \right)^{k_1} \dots \left(\frac{\partial y_1}{\partial x_n} \right)^{k_n},$$

is different from zero at the point $p \in S$.

Recall that around $p \in S$, the surface S is the level set $y_1^{-1}\{0\}$, hence, the gradient

$$(21) \quad Dy_1(p) = \left(\frac{\partial y_1}{\partial x_1} \Big|_p, \frac{\partial y_1}{\partial x_2} \Big|_p, \dots, \frac{\partial y_1}{\partial x_n} \Big|_p \right)$$

is a multiple of the unit vector $\nu_p := (\nu_1, \dots, \nu_n)$ normal to the surface S at p .

The *characteristic determinant* is the function $H : \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$(22) \quad H(x) = \det(m_{ij}(x)) = \det \left(\sum_{k_1+\dots+k_n=n_j} a_{ij}^{k_1 \dots k_n} x_1^{k_1} \dots x_n^{k_n} \right),$$

and the equation $H(x) = 0$ is called the *characteristic equation*.

Definition 2.2. The surface S is said to be *characteristic* at p if the characteristic equation is satisfied by the normal $\nu = (\nu_1, \dots, \nu_n)$ to S at p , and it will be simply called a *characteristic surface* if it is characteristic at every point.

Definition 2.3. The system (17) is called *elliptic* at p if for every surface S through p , S is non-characteristic, that is, the characteristic equation has no real solution other than the trivial, i.e., $H(x) = 0$ only for $x = 0$.

Remark On real characteristic curves it is not possible to eliminate at least one highest order derivative of u_j , hence, elliptic equations have no real characteristic curves.

3. MAXIMUM PRINCIPLE FOR THE LAPLACE EQUATION

This is one of the most important and fundamental results of partial differential equations of the elliptic type. There are many important properties of PDE's that follows if the maximum principle holds and the idea is the following: consider a smooth function $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$. The symmetric matrix of second order derivative has important information, for instance, one knows that if there is a local maximum at a point $x_0 \in \Omega$ then the Hessian matrix $D^2u(x_0)$ is negative definite, and the graph of u looks locally concave, while if x_0 is a local minimum the hessian matrix $D^2u(x_0)$ is positive definite, and the graph of u in this case looks locally convex. Since $\Delta u = \text{Tr}(D^2u)$, the condition $\Delta u > 0$ in Ω suggests that the graph of the function u looks (in some sense) convex.

Proposition 3.1. *If $\Delta u > 0$ in Ω , then u attains its maximum value at the boundary $\partial\Omega$.*

Proof. If there is a local maximum of $u \in C^2(\Omega)$ at x_0 in the interior of Ω we would have $\frac{\partial^2 u}{\partial x_i^2}(x_0) \leq 0$ for all $i = 1, 2, \dots, n$, which leads to a contradiction. \square

Proposition 3.2. *If $\Delta u \geq 0$ in Ω , then u attains its maximum value at the boundary $\partial\Omega$. Further more, for any ball $\overline{B_r(x_0)} \subset \Omega$ it holds that $u(x_0) \leq \max_{x \in \partial B_r(x_0)} u(x)$.*

Proof. On the contrary lets assume there is $r > 0$ such that $\overline{B_r(x_0)} \subset \Omega$ and $u(x_0) > m$, where $m := \max_{x \in \partial B_r(x_0)} u(x)$. Now define the function

$$(23) \quad U(x) := u(x) + \frac{\alpha}{r^2} |x - x_0|^2,$$

where $\alpha > 0$ is a constant to be determine later. Note that $U(x_0) = u(x_0)$. On the other hand if $|x - x_0| = r$ we have

$$(24) \quad U(x) = u(x) + \alpha < m + \alpha.$$

Choose any $0 < \alpha < u(x_0) - m$ and then for $|x - x_0| = r$ it holds

$$(25) \quad U(x) < u(x_0).$$

We also note

$$(26) \quad \Delta U(x) = \Delta u(x) + \frac{2\alpha n}{r^2} > 0,$$

which contradicts the previous proposition. \square

The following will be known as the mean value property (inequalities) for harmonic, subharmonic, superharmonic functions.

Proposition 3.3 ([3], T2.1). *Let $u \in C^2(\Omega)$. For any ball $B = B_R(y) \subset \subset \Omega$, we have the following:*

- *If u is harmonic in Ω , i.e., $\Delta u = 0$, then:*

$$(27) \quad u(y) = \frac{1}{n\omega_n R^{n-1}} \int_{\partial B} u ds.$$

$$(28) \quad u(y) = \frac{1}{\omega_n R^n} \int_B u dx.$$

- *If u is subharmonic in Ω , i.e., $\Delta u \geq 0$, then the equality should be replace with \leq .*
- *If u is superharmonic in Ω , i.e., $\Delta u \leq 0$, then the equality should be replace with \geq .*

Proof. Let $0 < \rho < R$.

Notice that for any $x \in \partial B_\rho(y)$ we have that the outer normal vector is

$$(29) \quad \nu(x) = \frac{x - y}{|x - y|} = \frac{x - y}{\rho},$$

and we can also show that

$$(30) \quad \frac{\partial u}{\partial \nu}(x) = \nabla u(x) \cdot \nu(x) = \frac{\partial u}{\partial \rho}(y + \rho \nu(x)).$$

This allows us to write

$$(31) \quad \int_{\partial B} \frac{\partial u}{\partial \nu}(x) dS_x = \int_{\partial B} \frac{\partial u}{\partial \rho}(y + \rho \nu(x)) dS_x.$$

Now, there is a relationship between the area element dS of ∂B_ρ and the area element $d\nu$ of the unit sphere \mathbb{S}^{n-1} , namely

$$dS = \rho^{n-1} d\nu.$$

Then we have the following sequence of equalities

$$\begin{aligned}
 0 &= \int_{B_\rho} \Delta u dx = \int_{\partial B_\rho} \frac{\partial u}{\partial \nu}(x) dS_x = \int_{\partial B_\rho} \frac{\partial u}{\partial \rho} u(y + \rho\nu(x)) dS_x \\
 &= \rho^{n-1} \int_{\mathbb{S}^{n-1}} \frac{\partial}{\partial \rho} u(y + \rho\nu) d\nu \\
 &= \rho^{n-1} \frac{\partial}{\partial \rho} \left(\int_{\mathbb{S}^{n-1}} u(y + \rho\nu) d\nu \right) \\
 &= \rho^{n-1} \frac{\partial}{\partial \rho} \left(\frac{\rho^{n-1}}{\rho^{n-1}} \int_{\mathbb{S}^{n-1}} u(y + \rho\nu) d\nu \right) \\
 &= \rho^{n-1} \frac{\partial}{\partial \rho} \left(\frac{1}{\rho^{n-1}} \int_{\partial B_\rho} u(x) dS_x \right)
 \end{aligned} \tag{32}$$

which implies that the quantity in brackets is constant (independent of $0 < \rho \leq R$), and in particular that

$$\rho^{1-n} \int_{\partial B_\rho} u(x) dS = R^{1-n} \int_{\partial B} u(x) dS. \tag{33}$$

and

$$\lim_{\rho \rightarrow 0} \rho^{1-n} \int_{\partial B_\rho} u(x) dS = \lim_{\rho \rightarrow 0} \int_{\mathbb{S}^{n-1}} u(y + \rho\nu) d\nu = \int_{\mathbb{S}^{n-1}} u(y) d\nu = u(y) n\omega_n. \tag{34}$$

The last two equations show that

$$u(y) = \frac{1}{n\omega_n R^{n-1}} \int_{\partial B} u(x) dS.$$

To obtain the solid form of the mean value property use the previous identity

$$u(y) \rho^{n-1} = \frac{1}{n\omega_n} \int_{\partial B_\rho} u(x) dS,$$

and integrate with respect to ρ from 0 to R . □

4. THE POISSON'S EQUATION IN DOMAINS OF \mathbb{R}^n

The most important example of an elliptic equation is the **Laplace equation** $\Delta u = 0$. The solutions of this equation are called **harmonic functions**. When the right hand side is different from zero, $\Delta u = f$, it is called **Poisson's equation**. The following points outline one path to establish the solvability of the Laplace and Poisson's equations using the so called Perron's method in bounded domains of the Euclidean space:

- First we will start with Poisson's equation in the unbounded \mathbb{R}^n by means of the Newton's kernel.
- After learning some properties of the Newton kernel and convolution with functions, we obtain the Green's representation formula. As an application we also obtain the Green's function in a ball for the Laplace equation.
- The solution of the Laplace equation in the ball is given explicitly by the Poisson's integral formula.
- Then we can show that in the ball, a solution of the Laplace equation, has certain regularity.
- To solve the Laplace equation in bounded domains Ω we make use of Perron's method for subharmonic functions.
- Then the solvability of the Dirichlet problem for Laplace equation regarding the boundary data is addressed.
- Poisson's equation in bounded domains will follow from Laplace equation, using suitable function in Newton's kernel, and Perron's method with the corresponding boundary data.
- We keep developing estimates for the solutions and establish Kellogg's theorem. This kind of a priori estimates will be further developed in the next section to establish existence of solutions for linear equations avoiding potential theory, i.e., by obtaining a priori estimates.

We are basically using the so called Potential theory in the ball, and then Perron's method to get a solution with certain regularity. This is also the underlying idea for proving the existence of a solution in the *viscosity* sense for fully nonlinear equations.

We start by considering the Poisson's equation in \mathbb{R}^n namely

$$\Delta u = f. \tag{35}$$

For compactly supported functions f , an explicit solution can be obtained using the Newton's kernel

$$(36) \quad \Gamma(x) = \begin{cases} c|x|^{2-n}, & n \geq 3 \\ -c \log |x|, & n = 2 \end{cases},$$

and then the solution is given by the convolution:

$$(37) \quad u(x) = (\Gamma * f)(x) := \int \Gamma(x-y)f(y)dy.$$

It is common to use $c = \frac{1}{2\pi}$ for $n = 2$ and $c = \frac{1}{n(2-n)\omega_n}$ for $n > 2$ in equation (36), where ω_n denotes the volume of the unit ball $B_1(0)$, and $\mathbb{S}^{n-1} = \partial B_1(0)$. We also consider the following identities:

$$(38) \quad n\omega_n R^n = n\text{Vol}(B_R) = \text{Area}(\mathbb{S}^{n-1})R^n = \text{Area}(\partial B_R)R.$$

Proposition 4.1. *Let $n > 2$ and suppose $y \in \mathbb{R}^n$ is fixed. Take $c = \frac{1}{n(2-n)\omega_n}$ in (36). If $x \neq y$ and writing $r^2 = |x-y|^2$, then the following identities hold:*

$$\begin{aligned} (1) \quad & \frac{\partial r}{\partial x_i} = \frac{x_i - y_i}{r}. \\ (2) \quad & \frac{\partial^2 r}{\partial x_j \partial x_i} = \frac{r^2 \delta_{ij} - (x_i - y_i)(x_j - y_j)}{r^3}. \\ (3) \quad & \frac{\partial \Gamma(x-y)}{\partial x_i} = \frac{r^{-n}}{n\omega_n} (x_i - y_i). \\ (4) \quad & \frac{\partial^2 \Gamma(x-y)}{\partial x_j \partial x_i} = \frac{r^{-n-2}}{n\omega_n} \{ \delta_{ij} r^2 - n(x_i - y_i)(x_j - y_j) \}. \\ (5) \quad & \frac{\partial^3 \Gamma(x-y)}{\partial x_k \partial x_j \partial x_i} = -\frac{(n+2)r^{-n-4}}{n\omega_n} (x_k - y_k) \{ \delta_{ij} r^2 - n(x_i - y_i)(x_j - y_j) \} \\ & + \frac{r^{-n-2}}{n\omega_n} \{ 2\delta_{ij}(x_k - y_k) - n\delta_{ik}(x_j - y_j) - n\delta_{jk}(x_i - y_i) \}. \\ (6) \quad & \Delta_x \Gamma(x-y) = 0. \end{aligned}$$

Proposition 4.2. *Let $n > 2$ and take $c = \frac{1}{n(2-n)\omega_n}$ in (36). Then $\Gamma(x)$ is integrable in the ball $B_R(0)$.*

Proof. If we write $x \in \mathbb{R}^n$ in polar coordinates, $x = r\xi$, where $r^2 = |x|^2$ and $\xi \in \mathbb{S}^{n-1}$, then the volume form dx of \mathbb{R}^n is expressed as $r^{n-1}drdS$, where dS is the volume form of \mathbb{S}^{n-1} . For any $R > 0$ fixed, it holds

$$\begin{aligned} (39) \quad \int_{B_R(0)} \Gamma(x)dx &= \frac{1}{n(2-n)\omega_n} \int_{B_R(0)} r^{2-n}dx \\ &= \frac{1}{n(2-n)\omega_n} \int_{\mathbb{S}^{n-1}} \int_0^R r^{2-n} r^{n-1} dr dS \\ &= \frac{1}{n(2-n)\omega_n} \int_{\mathbb{S}^{n-1}} \int_0^R r dr dS \\ &= \frac{R^2}{2(2-n)}. \end{aligned}$$

□

The following estimates will be useful, put $r = |x-y|$, then:

$$(40) \quad \left| \int_{\partial B_R(x)} \frac{\partial \Gamma(r)}{\partial x_i} dS_y \right| \leq C.$$

$$(41) \quad \left| \int_{\partial B_R(x)} \frac{\partial^2 \Gamma(r)}{\partial x_j \partial x_i} dS_y \right| \leq C R^{-1}.$$

$$(42) \quad \left| \int_{\partial B_R(x)} \frac{\partial^3 \Gamma(r)}{\partial x_k \partial x_j \partial x_i} dS_y \right| \leq C R^{-2}.$$

$$(43) \quad \left| \int_{B_R(x)} \frac{\partial^2 \Gamma(r)}{\partial x_j \partial x_i} r^\alpha dy \right| \leq \frac{C}{\alpha} R^\alpha.$$

Note that the convolution is commutative $\Gamma * f = f * \Gamma$, and if additionally f is smooth and has compact support, we have the following properties:

- (i) $\Gamma * \Delta f = f$,
- (ii) $\Delta(\Gamma * f) = f$.

Recall integration by parts over a domain Ω in \mathbb{R}^n for smooth function u and a vector field F , and the Green's theorem

$$(44) \quad \begin{aligned} \int_{\Omega} u \operatorname{div} F &= \int_{\Omega} \operatorname{div}(uF) - \int_{\Omega} F \cdot \nabla u, \\ &= \int_{\partial\Omega} uF \cdot \nu \, ds - \int_{\Omega} F \cdot \nabla u, \end{aligned}$$

where ν is the outward unit vector, ∇u is the gradient vector and \cdot denotes the usual inner product in \mathbb{R}^n . Then the identity $\Gamma * \Delta f = f$ follows after integrating by parts in the complement of a small neighbourhood of the origin and then taking the limit when the radius of this ball goes to zero. The second identity follows from the first one and noticing that we can move the Laplace operator Δ inside the integral with respect to the variable that does not appear as an argument of Γ .

Using integration by parts and using limit properties of the Newton kernel, one can obtain the *Green's representation formula*, for all $y \in \Omega$:

$$(45) \quad u(y) = \int_{\partial\Omega} \left(u \frac{\partial \Gamma}{\partial \nu}(x-y) - \frac{\partial u}{\partial \nu} \Gamma(x-y) \right) ds + \int_{\Omega} \Gamma(x-y) \Delta u \, dx.$$

Moreover, suppose $h \in C^1(\bar{\Omega}) \cap C^2(\Omega)$ is harmonic in Ω . $G = \Gamma + h$ and suppose $G = 0$ on $\partial\Omega$, then it is possible to obtain the Green's function of the first kind for the domain Ω :

$$(46) \quad u(y) = \int_{\partial\Omega} u \frac{\partial G}{\partial \nu} ds + \int_{\Omega} G \Delta u \, dx.$$

Green's function on the ball B_R with centre at the origin can be determined using the inversion through the sphere of radius R , given by $i(y) = \frac{R^2}{|y|^2} y$. Note that

$$(47) \quad \frac{|y|^2}{R^2} |x - i(y)|^2 = \frac{|x|^2 |y|^2}{R^2} - 2x \cdot y + R^2.$$

Then the Green function is given by

$$(48) \quad G(x, y) = \begin{cases} \Gamma(|x-y|) - \Gamma\left(\frac{|y|}{R} |x-i(y)|\right), & y \neq 0 \\ \Gamma(|x|) - \Gamma(R), & y = 0. \end{cases}$$

Let us consider now the following Dirichlet problem in the ball

$$(49) \quad \begin{cases} \Delta u(x) = 0, & x \in B \\ u(x) = \varphi(x), & x \in \partial B \end{cases},$$

The **Poisson's kernel** is given by the normal derivative of G on the boundary ∂B_R

$$(50) \quad \frac{\partial G}{\partial \nu} = \frac{R^2 - |y|^2}{n\omega_n R} \frac{1}{|x-y|^n}, \quad x \in \partial B_R.$$

There is an explicit solution to (49) that follows from the expression:

$$(51) \quad u(x) = \int_{\partial B_R} K(x, y) \varphi(y) \, ds_y + \int_{B_R} G(x, y) f(y) \, dy,$$

where K is the Poisson kernel and G is the green function of the ball, and moreover, whenever φ is continuous in ∂B_R , then $u(x) \in C^0(\bar{B}_R) \cap C^2(B_R)$.

Theorem 4.3 ([3], T2.6). *Let $B = B_R(0)$ and φ a continuous function on ∂B . Then the function u given by*

$$(52) \quad u(x) = \begin{cases} \frac{R^2 - |x|^2}{n\omega_n R} \int_{\partial B} \frac{\varphi(y)}{|x-y|^n} \, ds_y, & x \in B \\ \varphi(x), & x \in \partial B \end{cases},$$

is harmonic and $u \in C^0(\bar{B}) \cap C^2(B)$.

Poisson's kernel is smooth for all $x \in B$, then the smoothness of u will follow from the fact that we can differentiate under the integral. This is a consequence of the Lebesgue' Dominated Convergence Theorem in measure theory: Let (Ω, μ) a measure space, and $X \subseteq \mathbb{R}$ open. Consider a function $f : X \times \Omega \rightarrow \mathbb{R}$, such that for every $x \in X$, the function $f(x, \cdot) : \Omega \rightarrow \mathbb{R}$ is a Lebesgue-integrable function; for all $x \in X$ the derivative $\partial_x f$ exists for almost all $\omega \in \Omega$; There is a Lebesgue integrable function $g : \Omega \rightarrow \mathbb{R}$ such that $|\partial_x f(x, \omega)| \leq g(\omega)$, for all $x \in X$ and for almost every $\omega \in \Omega$. Then applying the Dominated Convergence Theorem, for all $x \in X$ we have:

$$(53) \quad \frac{d}{dx} \int_{\Omega} f(x, \omega) \, d\mu = \int_{\Omega} \partial_x f(x, \omega) \, d\mu.$$

4.1. Perron's Method. Let $\Omega \in \mathbb{R}^n$ a domain. $u \in C^0(\Omega)$ is called **subharmonic** if its values are less or equal than the values of any harmonic function with bigger boundary data than u , all when restricted to any properly contained ball in Ω , more precisely, if for every ball B such that $\bar{B} \subset \Omega$ and every function h harmonic in B such that $u \leq h$ on ∂B then we also have $u \leq h$ in B . The definition of **superharmonic** functions follows by replacing \leq by \geq . Now we list some properties of subharmonic functions

- If u is subharmonic in Ω then it satisfies the strong maximum principle (if maximum is attained in the interior then it is constant).
- When comparing a superharmonic function v and a subharmonic function u in the a bounded domain Ω such that $v \geq u$ on $\partial\Omega$ then if $v \neq u$ we have $v > u$ in all Ω .
- Let u be subharmonic in Ω and B such that $\bar{B} \subset \Omega$. By the Poisson's integral of u in ∂B we obtain a harmonic function \bar{u} in B such that $\bar{u} = u$ on ∂B . From this, we can construct the following function U , which is subharmonic in Ω given by

$$(54) \quad U(x) = \begin{cases} \bar{u}(x), & x \in B \\ u(x), & x \in \Omega \setminus B. \end{cases}$$

- Let $\{u_1, \dots, u_m\}$ be a finite collection of subharmonic functions in Ω . Then $u(x) = \max_i \{u_i(x)\}$ is subharmonic.

The corresponding results for superharmonic functions follow from replacing u by $-u$ in each property.

Let Ω be a bounded domain and φ a function in Ω that is bounded in $\partial\Omega$. Then a $C^0(\bar{\Omega})$ function u that is subharmonic in Ω is called **subfunction** relative to φ , if $u \leq \varphi$ on $\partial\Omega$. We define **superfunction** in a similar way. Note that constant functions with value less or equal than $\inf_{\partial\Omega} \varphi$ are subfunctions. Also note that by the maximum principle every subfunction is less or equal than every superfunction.

Theorem 4.4 (Perron's Method ([3], T2.12)). *Let S_φ the set of subfunctions relative to the function $\varphi : \Omega \rightarrow \mathbb{R}$, where Ω is a bounded domain of \mathbb{R}^n and φ is bounded on $\partial\Omega$. Then the function*

$$(55) \quad u(x) = \sup_{v \in S_\varphi} v(x),$$

is harmonic in Ω .

One way to prove Perron's methods is via Harnack's convergence theorem using Harnack's inequality (see at the end of this section), by showing that u can be approximated by harmonic functions. Another way to prove it is by using interior derivative estimates for harmonic functions, which imply the equicontinuity on compact subdomains of *the second order derivatives* of any (uniformly) bounded collection of harmonic functions. Consequently by Arzela's theorem we have that any bounded sequence of harmonic functions forms a normal family. In order to see this, it is worth analysing the following example:

Example 4.5 ([3], T2.10). *From the mean value property and divergence theorems it follows*

$$(56) \quad D_i u(x) = \frac{1}{\omega_n R^n} \int_B D_i u \, dy = \frac{1}{\omega_n R^n} \int_B \operatorname{div}(u e_i) \, dy = \frac{1}{\omega_n R^n} \int_{\partial B} u e_i \cdot \nu \, ds = \frac{1}{\omega_n R^n} \int_{\partial B} u \nu_i \, ds.$$

One can show that if u is harmonic in a domain Ω and if Ω' is any compact subset of Ω and for any multi-index r with $d = \operatorname{dist}(\Omega', \partial\Omega)$ then following inequality holds

$$(57) \quad \sup_{\Omega'} |D^r u| \leq \left(\frac{n|r|}{d} \right)^{|r|} \sup_{\Omega} |u|.$$

Note that for any open convex subset of \mathbb{R}^n we have the estimate

$$(58) \quad |u(x) - u(y)| \leq |Du(z)| |x - y|.$$

*Whenever the partial derivatives of u are bounded, u is Lipschitz continuous. Moreover, any bounded sequence of harmonic functions on a domain Ω contains a subsequence converging uniformly on compact subdomains of Ω to a continuous function. To show that this limit function is also a harmonic function, take any ball B contained in Ω , and for each element of the sequence, write the mean value property. Then the uniform convergence will imply that the mean value property is also satisfied by the limit function. Then conclude that **any bounded sequence of harmonic functions in a domain Ω contains a subsequence converging uniformly on compact subdomains of Ω to a harmonic function.** ([3], T2.11).*

Perron's method then ensures the existence of a solution to the equation $\Delta u = 0$ in a bounded domain Ω of \mathbb{R}^n , although it is sensible to expect that $u \leq \varphi$ on $\partial\Omega$. In the case that u is a solution of the *solvable Dirichlet problem* $\Delta u = 0$ in Ω , $u = \varphi$ on $\partial\Omega$, then this solution coincide with the solution given by Perron's method. However *the solvability of the Dirichlet problem depends on the geometric properties of the boundary $\partial\Omega$* . For each $z \in \partial\Omega$, we define the **barrier** at z relative to Ω , denoted by w_z , to be a $C^0(\bar{\Omega})$ function such that w_z is superharmonic in Ω , $w_z > 0$ in $\bar{\Omega} \setminus \{z\}$ and $w_z(z) = 0$. A point $z \in \partial\Omega$ is called **regular** if there exists a barrier at z . It is at regular boundary points and for φ continuous at these, that the solution u in Perron's method, matches with the boundary data of the Dirichlet problem, i.e., $\lim_{x \rightarrow z} u(x) = \varphi(z)$.

Theorem 4.6 ([3], T2.14). *The classical Dirichlet problem with continuous boundary data is solvable if and only if the boundary points are all regular.*

The remaining problem is the characterisation of domains whose boundary points are all regular. In the case of $n = 2$, the Dirichlet problem is solvable if every component of the complement of the domain Ω has more than one element. Examples of this are domains bounded by a finite number of simple closed curves. In general dimension, when the boundary of Ω is C^2 then all boundary points are regular.

Example 4.7 (Exterior sphere condition and barrier). *Let $q \in \partial\Omega$ and let B_r such that $\bar{B}_r \cap \bar{\Omega} = \{q\}$. Then we can consider as barrier the following harmonic function*

$$(59) \quad w_q(p) = \begin{cases} \frac{1}{r^{n-2}} - \frac{1}{d^{n-2}}, & n > 2, \\ \ln\left(\frac{d}{r}\right), & n = 2. \end{cases}$$

where d is the distance of p to the centre of B_r .

Now we turn to the Poisson's equation $\Delta u = f$ and state the existence result for the corresponding Dirichlet problem in a domain $\Omega \subset \mathbb{R}^n$. First we note that if f is continuous, then the **Newton potential** $\Gamma * f$ is not necessarily twice differentiable. Notice as well that the solution is not unique, since adding a harmonic function on Ω gives another solution, nevertheless, when adding boundary conditions, one can obtain existence and uniqueness results.

Example 4.8. *In $\Omega = \{(x_1, x_2) \in \mathbb{R}^2 \mid x_1, x_2 > 0\}$, let $u(x_1, x_2) = x_1 x_2 \log(x_1 + x_2)$ and $f(x_1, x_2) = 2\left(1 - \frac{x_1 x_2}{(x_1 + x_2)^2}\right)$. Then $\Delta u = f$, however $f \in L^\infty(\Omega)$ but $u \notin C^{1,1}(\Omega)$, since $u_{x_1 x_2}$ is not bounded in Ω .*

Example 4.9. *For the Newton potential $\Gamma * f$ to be twice differentiable, it is necessary to assume f is Hölder continuous. In this case, one can obtain a $C^2(B)$ bound for a solution of the problem $\Delta u = f$ in $\Omega = B_d(x) = B$. Notice that by integration by parts, the solution is written as*

$$(60) \quad u(x) = \int_{\partial\Omega} \left(\Gamma(x-y) \frac{\partial u(y)}{\partial \nu} - u(y) \frac{\partial \Gamma(x-y)}{\partial \nu} \right) dS_y - \int_{\Omega} \Gamma(x-y) f(y) dy.$$

Using properties of the Newtonian potential we have

$$(61) \quad \begin{aligned} v(x) &= - \int_{\Omega} \Gamma(x-y) f(y) dy, \\ \frac{\partial v(x)}{\partial x_i} &= - \int_{\Omega} \frac{\partial \Gamma(x-y)}{\partial x_i} f(y) dy, \\ \frac{\partial^2 v(x)}{\partial x_j \partial x_i} &= f(x) \int_{\partial\Omega} \frac{\partial \Gamma(x-y)}{\partial x_j} \nu_i dS_y - \int_{\Omega} \frac{\partial^2 \Gamma(x-y)}{\partial x_j \partial x_i} (f(y) - f(x)) dy. \end{aligned}$$

Then it is possible to get an estimate for the second derivatives of u . Since

$$(62) \quad \begin{aligned} \frac{\partial^2 u(x)}{\partial x_j \partial x_i} &= \int_{\partial\Omega} \frac{\partial^2 \Gamma(x-y)}{\partial x_j \partial x_i} \frac{\partial u(y)}{\partial \nu} dS_y - \int_{\partial\Omega} u(y) \frac{\partial}{\partial \nu} \frac{\partial^2 \Gamma(x-y)}{\partial x_j \partial x_i} dS_y \\ &\quad + f(x) \int_{\partial\Omega} \frac{\partial \Gamma(x-y)}{\partial x_j} \nu_i dS_y - \int_{\Omega} \frac{\partial^2 \Gamma(x-y)}{\partial x_j \partial x_i} (f(y) - f(x)) dy, \end{aligned}$$

then the following inequality holds

$$(63) \quad |D^2 u(x)| \leq C \left(d^{-2} \max_{\partial B_d(x)} |u| + d^{-1} \max_{\partial B_d(x)} |Du| + |f(x)| + d^\alpha |f|_\alpha \right),$$

where $C = C(n, \alpha)$.

Theorem 4.10 ([3], T4.3). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and suppose that every boundary point is regular. Assume that f is bounded and locally Hölder continuous in Ω . Then for any continuous boundary values φ , the Dirichlet problem*

$$(64) \quad \begin{cases} \Delta u = f & \text{in } \Omega \\ u = \varphi & \text{on } \partial\Omega \end{cases},$$

has a unique solution.

Proof. Note that if w is the Newtonian potential of f and if we consider $v = u - w$, the last problem is equivalent to Laplace's equation $\Delta v = 0$ in Ω and boundary values $\varphi - w$. \square

4.2. Interior estimates. The following is to address the question of the regularity of the solution after this limiting process. One can prove (see [3], L4.4) that if $B_1 \subset B_2$ are two concentric balls, $f \in C^\alpha(\bar{B}_2)$ where $0 \leq \alpha \leq 1$ then $w := \Gamma * f$ is $C^{2,\alpha}$ in \bar{B}_1 . One can improve this Hölder estimate to solutions of $\Delta u = f$, where u and f have compact support, so u belongs to $C_0^2(\mathbb{R}^n)$ and f in $C_0^\alpha(\mathbb{R}^n)$, implying that $u \in C_0^{2,\alpha}(\mathbb{R}^n)$ and moreover:

Theorem 4.11 ([3], T4.6). *Let $\Omega \subset \mathbb{R}^n$ a domain and suppose $u \in C^2(\Omega)$ and $f \in C^\alpha(\Omega)$ are such that $\Delta u = f$ in Ω . Then $u \in C^{2,\alpha}(\Omega)$. Moreover for any concentric balls $B_R \subset B_{2R} \subset \subset \Omega$ one can estimate*

$$(65) \quad |u|'_{C^{2,\alpha}(B_R)} \leq C(|u|_{C^0(B_{2R})} + R^2|f|'_{C^{0,\alpha}(B_{2R})}),$$

where $C = C(n, \alpha)$.

Last result gives the interior $C^{2,\alpha}$ regularity of solutions inside properly contained balls. See an application of the interior regularity for linear operators below. As we have noted earlier, an immediate consequence of this interior estimate is the equicontinuity on compact subdomains of the *second derivatives* of any bounded family of solutions of the equation $\Delta u = f$. In the case of harmonic functions, interior estimates of derivatives and Arzela's Theorem is used to establish in [3], T2.11 that any bounded family of harmonic functions is a *normal family*. For the Poisson's equation then (65) and Arzela's theorem imply that any bounded family of solution to $\Delta u = f$ in Ω , with $f \in C^\alpha(\Omega)$, contains a subsequent converging uniformly on compact subdomain to a solution. This compactness result gives the existence result for Poisson's equation:

Theorem 4.12 ([3]T4.9). *Let B a ball in \mathbb{R}^n and $f \in C^\alpha(B)$ such that*

$$(66) \quad \sup_{x \in B} d_x^{2-\beta} |f(x)| \leq N < \infty, \quad \text{for some } 0 < \beta < 1.$$

Then there is a unique solution $u \in C^0(\bar{B}) \cap C^2(B)$, of the Dirichlet problem

$$(67) \quad \begin{cases} \Delta u = f & \text{in } B \\ u = 0 & \text{on } \partial B, \end{cases}$$

Moreover the following inequality holds

$$(68) \quad \sup_{x \in B_R} d_x^{-\beta} |u(x)| \leq CN, \quad \text{where the constant } C = C(\beta).$$

On the other hand, from Theorem 4.12 one can obtain a similar estimate replacing general domains Ω (not necessarily bounded, [3], T4.8). Moreover this estimate is also applicable to the intersection of the domain Ω and the upper half space \mathbb{R}_+^n ([3], T4.11).

4.3. Regularity of solutions to the boundary value problem: Kellogg's Theorem. The next result is about the regularity of the solutions of the Dirichlet problem in the ball and up to the boundary. It states that when a solution is in $C^0(\bar{B}_R) \cap C^2(B_R)$, then this is also in $C^{2,\alpha}(\bar{B})$. The result follows from ([3], T4.11) applied to the Kelvin transformation.

Theorem 4.13 (Kellogg's Theorem in the Ball ([3], T4.13, C4.14)). *Let $f \in C^\alpha(\bar{B})$ and $\varphi \in C^{2,\alpha}(\bar{B})$, and let $u \in C^0(\bar{B}) \cap C^2(B)$ a solution of the Dirichlet problem*

$$(69) \quad \begin{cases} \Delta u = f & \text{in } B \\ u = \varphi & \text{on } \partial B, \end{cases}$$

where $B \subset \mathbb{R}^n$ is a ball. Then $u \in C^{2,\alpha}(\bar{B})$, and the solution is unique.

Kellogg's theorem is valid for more general domains $\Omega \subset \mathbb{R}^n$, and will be discussed in the following section.

Remark. Regarding the regularity of the solutions we note that it is possible to prove using mollifiers that if u is a C^2 solution of the Laplace's equation, then u is smooth. Even more, if u is assumed to be continuous in the domain U and satisfies the mean value property for each ball inside the domain, then u is smooth, since its mollification is the same function. Recall that a mollifier is a smooth function μ with the property that it can make sharp pieces of a given function to be *mollified* or smooth-out, after convolution. Some facts about mollifiers are: a) Re-scaling in ϵ -balls, the convolution of an ϵ -mollifier μ_ϵ with a locally integrable function is smooth. b) Almost everywhere, the point-wise convergence of $(\mu_\epsilon * f) \rightarrow f$ as $\epsilon \rightarrow 0$ holds. c) If f is continuous, the convergence is uniformly on compact subsets of the domain. d) if f is in L_{loc}^p , $1 \leq p < \infty$, the convergence is also in L_{loc}^p sense.

4.4. Harnack's Inequality. (See [5]) Suppose that u is harmonic and non-negative in the Ball $B = B_R(0)$. If $x \in B$ and $y \in \partial B$ then we have

$$R - |x| = ||x| - |y|| \leq |x - y| \leq |x| + |y| = R + |x|,$$

this implies

$$\frac{R^2 - |x|^2}{nw_n R(R - |x|)^n} \geq \frac{R^2 - |x|^2}{nw_n R|y - x|^n} \geq \frac{R^2 - |x|^2}{nw_n R(R + |x|)^n},$$

or equivalently

$$\frac{R + |x|}{R - |x|} \frac{1}{nw_n R(R - |x|)^{n-2}} \geq \frac{R^2 - |x|^2}{nw_n R|y - x|^n} \geq \frac{R - |x|}{R + |x|} \frac{1}{nw_n R(R + |x|)^{n-2}},$$

after substituting in the Piosson's Integral Formula and a limit argument one can get the inequalities

$$\left(\frac{R}{R - |x|}\right)^{n-2} \frac{R + |x|}{R - |x|} u(0) \geq u(x) \geq \left(\frac{R}{R + |x|}\right)^{n-2} \frac{R - |x|}{R + |x|} u(0).$$

One consequence Harnack's inequality is that any harmonic function defined for all \mathbb{R}^n and bounded from below is identically constant, this follows easily by adding a constant if necessary to have the function to be positive and by taking the limit as R tends to infinity.

Also it is possible to prove that if $\{u_n\}$ is a monotone increasing sequence of harmonic functions bounded from above at a point $p \in \Omega$, then in each compact subset of Ω , $\{u_n\}$ converges uniformly to a harmonic function.

A generalisation of this convergence result for uniform limit of solutions to a linear second order PDE will be a consequence of the Schauder's interior estimates, to be discuss in the following section.

5. LINEAR SECOND ORDER PARTIAL DIFFERENTIAL EQUATIONS: SCHAUDER THEORY.

There exists a complete theory for linear equations relying in a priori estimates. These are inequalities on the values of solutions and its derivatives, before actually guarantee its existence.

By solving the Poisson's equation $\Delta u = f$ in a domain $\Omega \subset \mathbb{R}^n$ it is possible to show that for any subset $\Omega' \subset \subset \Omega$, if $u \in C^2(\Omega)$ is a solution of the equation, then

$$(70) \quad \|u\|_{C^{2,\alpha}(\bar{\Omega}')} \leq C(\|u\|_{C^0(\Omega)} + \|f\|_{C^\alpha(\bar{\Omega})}),$$

with $C = C(\alpha, n, d(\Omega', \partial\Omega))$, where $\alpha \in (0, 1)$ and $d(\Omega', \partial\Omega)$ is the distance between the sets. This estimate can be turned into a global estimate for solutions with sufficiently smooth boundary values, and with boundary $\partial\Omega$ sufficiently smooth.

Schauder theory consists in obtaining the same inequality for any $C^{2,\alpha}$ solution of $Lu = f$, where now the constant C depends additionally on bounds of the coefficients in Hölder space sense and also on the minimum and maximum eigenvalues of the coefficient matrix (a^{ij}) in Ω .

These estimates are used to apply the Continuity method and obtain the existence of a solution for the Dirichlet from perturbations of the Poisson's equation.

For solutions of elliptic linear equations in bounded domains, the maximum principle gives the following point wise estimate

Theorem 5.1 ([3], T3.7). *Let $Lu = f$ in a bounded domain $\Omega \subset \mathbb{R}^n$ where L is elliptic, $c \leq 0$ and $u \in C^0(\bar{\Omega}) \cap C^2(\Omega)$ is a solution. Then*

$$(71) \quad |u|_{C^0(\Omega)} \leq \sup_{\partial\Omega} |u| + C \sup_{\Omega} \frac{|f|}{\lambda},$$

where $C = C(\text{diam } \Omega, \beta = \sup |b|/\lambda)$.

Theorem 5.2 (Schauder's estimate, ([3], T6.6)). *Let $\Omega \subset \mathbb{R}^n$ be a $C^{2,\alpha}$ domain and assume $u \in C^{2,\alpha}(\bar{\Omega})$ is a solution of $Lu = f$ in Ω and $u = \varphi$ on $\partial\Omega$, where $\varphi \in C^{2,\alpha}(\bar{\Omega})$, $f \in C^\alpha(\bar{\Omega})$, L is uniformly elliptic. Also assume that there are positive constants λ, Λ such that for $a^{ij}, b^i, c \in C^\alpha(\bar{\Omega})$ we have*

$$(72) \quad a^{ij}(x) \xi_i \xi_j \geq \lambda |\xi|^2, \quad x \in \Omega, \xi \in \mathbb{R}^n,$$

and

$$(73) \quad |a^{ij}|_{C^\alpha(\Omega)}, |b^i|_{C^\alpha(\Omega)}, |c|_{C^\alpha(\Omega)} \leq \Lambda.$$

Then there is a constant $C = C(n, \alpha, \lambda, \Lambda, \Omega)$ such that the following inequality holds

$$(74) \quad |u|_{C^{2,\alpha}(\Omega)} \leq C(|u|_{C^0(\Omega)} + |\varphi|_{C^{2,\alpha}(\bar{\Omega})} + |f|_{C^\alpha(\bar{\Omega})}).$$

Proof. Interior C^2 estimates. To obtain these estimates we proceed as follows:

Let d_x the distance from x to the boundary $\partial\Omega$, and let $d_{xy} = \min\{d_x, d_y\}$. Then for the equation in Ω

$$(75) \quad Lu = a^{ij}u_{ij} + b^i u_i + cu = f,$$

define

$$(76) \quad \begin{aligned} M_1 &:= \sup_{x \in \Omega} d_x |Du(x)|, \\ M_2 &:= \sup_{x \in \Omega} d_x^2 |D^2u(x)|, \\ M_{j+\alpha} &= \sup_{x, y \in \Omega} d_{xy}^{j+\alpha} \frac{|u(x) - u(y)|}{|x - y|^\alpha}. \end{aligned}$$

And for any function $f : \Omega \rightarrow \mathbb{R}$

$$(77) \quad H_{p, \Omega}[u] := \sup_{x \in \Omega} \frac{|u(x) - u(p)|}{|x - p|^\alpha}.$$

Then by definition there is a $p \in \Omega$ such that for $\epsilon = \frac{1}{2}M_2$ there is a second partial derivative of u at p such that

$$(78) \quad d_p^2 \left| \frac{\partial^2 u(p)}{\partial x_i \partial x_j} \right| > \frac{1}{2}M_2.$$

Consider a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$, and put $E = T(\Omega)$. Then for every $y \in E$, by writing $v(y) = (u \circ T^{-1})(y) = u(x)$, the equation (79) induces an equation in E

$$(79) \quad Lv = A^{ij}v_{ij} + B^i v_i + Cv = g.$$

Note that

$$(80) \quad \begin{aligned} A^{ij} &= a^{rs} \frac{\partial y_i}{\partial x_r} \frac{\partial y_j}{\partial x_s}, \\ B^i &= b^r \frac{\partial y_i}{\partial x_r}, \\ C &= c \circ T^{-1}, \\ g &= f \circ T^{-1}. \end{aligned}$$

Among the linear Transformations T we choose the one such that, for $q = T(p)$, it holds

$$(81) \quad A^{ij}(q) = \delta^{ij}.$$

From the ellipticity of (79) and since the coefficients are in C^α there are constant K and m such that

$$(82) \quad |A^{ij}|_\alpha, |B^i|_\alpha, |C|_\alpha, |g|_\alpha \leq K, \quad A^{ij}\xi_i\xi_j \geq m|\xi|^2.$$

For all $y \in E$, denote the distance from y to the boundary of E by e_y . Also consider B , the ball with centre at $q = T(p)$ and radius $d = \lambda e_q$, where $0 < \lambda < \frac{1}{2}$ is a constant to be determined later.

There is a constant $K_2 = K_2(K, m)$ such that

- Distance is stretched (or contracted) at most by a factor K_2 (or K_2^{-1}): $e_y = e_{tx} \leq K_2 d_x$.
- $|g|^* \leq K_2 |f|^*$
- $e_y |Dv(y)| \leq K_2 M_1$
- $e_y^2 |D^2v(y)| \leq K_2 M_2$
- $\sup_B |v| \leq M_0$
- $\sup_B |Dv| \leq \frac{K_2 M_1}{(1-\lambda)e_q}$.
- $\sup_B |D^2v| \leq \frac{K_2 M_2}{(1-\lambda)^2 e_q^2}$.
- $H_{q, B}[g] \leq (1-\lambda)^{-(2+\alpha)} |g|^*$
- $H_{q, B}[v] \leq \frac{K_2 M_1}{(1-\lambda)e_q} (\lambda e_q)^{1-\alpha}$.
- $H_{q, B}[Dv] \leq \frac{K_2 M_2}{(1-\lambda)^2 e_q^2} (\lambda e_q)^{1-\alpha}$.

From (78) it follows there exists partial derivatives in the y variables such that

$$(83) \quad e_p^2 \left| \frac{\partial^2 v(q)}{\partial x_i \partial x_j} \right| > K_3 M_2.$$

Now in E , consider the following Poisson's equation

$$(84) \quad \Delta v = (\delta^{ij} - A^{ij})v_{ij} - B^i v_i - Cv + g = (\Delta - L)v + g =: F.$$

Using the estimate (63)

$$(85) \quad |D^2v(q)| \leq C \left(d^{-2} \max_{\partial B_d(q)} |v| + d^{-1} \max_{\partial B_d(q)} |Dv| + |F(q)| + d^\alpha |F|_\alpha \right).$$

One can show that

$$(86) \quad d^{-2} \max_{\partial B_d(q)} |v| + d^{-1} \max_{\partial B_d(q)} |Dv| + |F(q)| \leq e_q^{-2} (K_6 \lambda^{-2} M_0 + K_6 \lambda^{-1} M_1 + |f|^*),$$

and

$$(87) \quad \begin{aligned} d^\alpha H_{p,B}[F] &\leq \lambda^\alpha e_q^\alpha K \left\{ \max_B |D^2 v| + \max_B |Dv| + H_{q,B}[Dv] + \max_B |v| + H_{q,B}[v] \right\} + \lambda^\alpha e_q^\alpha H_{q,B}[b] \\ &\leq \lambda^\alpha e_q^{-2} K_8 \{M_2 + M_1 + M_0 + |f|^*\}. \end{aligned}$$

Then we combine the inequalities and we have

$$(88) \quad e_q^2 |D^2 v(q)| \leq K_9 \{ \lambda^\alpha M_2 + \lambda^{-1} M_1 + \lambda^{-2} M_0 + |f|^* \}.$$

and hence

$$(89) \quad M_2 \leq K_{10} \{ \lambda^\alpha M_2 + \lambda^{-1} M_1 + \lambda^{-2} M_0 + |f|^* \}.$$

In order to apply a bootstrap argument we need an auxiliary result. Let u be a twice continuously differentiable function in the domain Ω . Let $0 < \mu < 1$. Then the following inequality holds:

$$(90) \quad M_1 \leq \frac{\mu M_2}{(1-\mu)^2} + \frac{M_0}{\mu}.$$

When applying this lemma to (89), one gets

$$(91) \quad M_2 \leq K_{10} \{ \lambda^\alpha M_2 + \lambda^{-1} \mu M_2 (1-\mu)^{-2} + (\lambda^{-1} \mu^{-1} + \lambda^{-2}) M_0 + |f|^* \}.$$

The choosing $\mu = \lambda^{1+\alpha}$, $\lambda < \frac{1}{2}$ and even more, choosing λ sufficiently small one gets

$$(92) \quad M_2 \leq K_{12} (M_0 + |f|^*).$$

Applying now the lemma again, we obtain

$$(93) \quad M_1 \leq K_{12} (M_0 + |f|^*).$$

Then we have the desired C^2 bound for u . □

From this a priori estimate then we can start with a solution for the Poisson's equation $\Delta u = f$ and then trace a solution for $Lu = f$ through solutions of a family of equations joining $\Delta u = f$ with $Lu = f$. This idea is stated in the following theorem.

Theorem 5.3 (Continuity Method ([3], T5.2)). *Let B be a Banach space, V a normed linear space, and $L_0, L_1 : B \rightarrow V$ two bounded linear operators. Define for $0 \leq t \leq 1$ the family*

$$(94) \quad L_t = (1-t)L_0 + tL_1,$$

and assume there exists a constant C such that

$$(95) \quad |x|_B \leq C |L_t x|_V, \quad 0 \leq t \leq 1.$$

Then L_1 is surjective if and only if L_0 is surjective.

Proof. Suppose that L_t is surjective, then by (95), the map L_t is invertible and such that $|L_t^{-1}| < C$, with C independent of t . Now consider $s \in [0, 1]$, since

$$(96) \quad \begin{aligned} L_t &= L_0 + t(L_1 - L_0) \\ L_s &= L_0 + s(L_1 - L_0), \end{aligned}$$

after subtraction we get for any $x \in B$

$$(97) \quad L_t x = L_s x + (s-t)(L_0 - L_1)x.$$

We will find conditions to ensure that L_s is onto. Let $y \in V$, and we want to solve $L_s x = y$. This is equivalent to show that

$$(98) \quad L_t x = y + (s-t)(L_0 - L_1)x,$$

which in turn is equivalent to

$$(99) \quad x = L_t^{-1} y + (s-t)L_t^{-1}(L_0 - L_1)x.$$

Now define

$$(100) \quad Tx := L_t^{-1} y + (s-t)L_t^{-1}(L_0 - L_1)x,$$

and notice that $T : X \rightarrow X$ is a contraction map whenever $|s-t| < \delta := [C(|L_0| + |L_1|)]^{-1}$.

Then the equation $Tx = x$ has a unique solution, which implies L_s is onto for $s \in [0, 1]$ with $|s - t| < \delta$. Then we only have to divide the interval $[0, 1]$ into subintervals of length less than δ and using the hypothesis that L_0 is surjective for a start. \square

First we are going to establish the existence for the Dirichlet problem in the case where Ω is the ball.

Proposition 5.4 ([3], C6.9). *Let B a ball in \mathbb{R}^n and L strictly elliptic in B , with coefficients in $C^\alpha(\bar{B})$ and $c \leq 0$. Then if $f \in C^\alpha(\bar{B})$ and $\varphi \in C^{2,\alpha}(\bar{B})$, there is a unique solution $u \in C^{2,\alpha}(\bar{B})$ of the boundary value problem $Lu = f$ in B , $u = \varphi$ on ∂B .*

Proof. Apply Theorem 5.3 as indicated in Theorem 5.7 below, replacing Ω with the ball B , and using Kellogg's theorem on the ball (Theorem 4.13). \square

Now we are going to establish the Perron's method for linear equations. We introduce the concept of **subsolution** of $Lu = f$, namely a function $u \in C^0(\Omega)$ such that for every ball B properly contained in Ω and every solution v of $Lv = f$ in B , if $u \leq v$ on ∂B then $u \leq v$ in B . By changing \leq by \geq , we say that $u \in C^0(\Omega)$ is a **supersolution** of $Lu = f$. When $f \in C^\alpha(\Omega)$ and also the coefficients of L are Hölder continuous, $c \leq 0$ and such that the maximum principle holds, then we have the following properties:

- The function $u \in C^2(\Omega)$ is a subsolution if and only if $Lu \geq f$.
- Let Ω be bounded, u and v a subsolution and a supersolution in Ω respectively. If $v \geq u$ on ∂B then either $v \geq u$ in Ω or they are equal.
- Let B a ball such that $\bar{B} \subset \Omega$ and u a subsolution in Ω . Let \bar{u} the solution to the Dirichlet problem in the ball $L\bar{u} = f$ in B such that $\bar{u} = u$ on ∂B . Then we can obtain a subsolution U in Ω given by

$$U(x) = \begin{cases} \bar{u}(x) & x \in B \\ u(x) & x \in \Omega \setminus B \end{cases}.$$

- Let u_1, \dots, u_k be subsolutions in Ω . Then the function $u(x) = \max_i \{u_i(x)\}$ is a subsolution in Ω .

Let Ω be bounded and φ a bounded function on $\partial\Omega$. We say that $u \in C^0(\bar{\Omega})$ is a **subfunction** relative to φ if u is a subsolution in Ω and $u \leq \varphi$ on $\partial\Omega$. We say that u is a **supersolution** relative to φ if it is a superfunction relative to φ in Ω and $u \geq \varphi$ on $\partial\Omega$. Define S_φ to be the set of all subfunctions in Ω relative to φ . In the case L is strictly elliptic in Ω and f and the coefficients of L are bounded, then S_φ is non-empty and bounded from above.

Theorem 5.5 (Perron's process, ([3], T6.11)). *Let Ω a bounded domain, L a strictly elliptic operator with $c \leq 0$ and coefficients in $C^\alpha(\Omega)$, the function $f \in C^\alpha(\Omega)$, and φ a bounded function on $\partial\Omega$. Then the function $u(x) = \sup_{v \in S_\varphi} v(x)$ belongs to $C^{2,\alpha}(\Omega)$ and is such that $Lu = f$ in Ω whenever u is bounded.*

Next we analyse the conditions that will make this solution to assume given boundary values. Let φ bounded on $\partial\Omega$ and continuous at $x_0 \in \partial\Omega$. The sequence of functions $\{w_i^+\}$ and $\{w_i^-\}$ are called respectively **upper barrier** and **lower barrier** in Ω relative to L, f and φ at the point $x_0 \in \Omega$ if each w_i^+ is a superfunction relative to φ in Ω , w_i^- is a subfunction relative to φ in Ω , and $\lim_{i \rightarrow \infty} w_i^\pm(x_0) = \varphi(x_0)$. If at a point there exist both upper and lower barrier, then we say simply that there is a barrier at that point.

Theorem 5.6 ([3], L6.12). *For a bounded function φ on $\partial\Omega$ continuous at $x_0 \in \partial\Omega$, the solution of the Dirichlet problem $Lu = f$ in Ω given by the Perron's process, satisfy the boundary condition $\lim_{x \rightarrow x_0} u(x) = \varphi(x_0)$ if there exist a barrier at x_0 .*

Again the question of what domains admit a barrier is of interest. In particular, any C^2 domain, and domains satisfying an exterior sphere condition at every point on the boundary, have a barriers ([3], T6.13).

Now we get the general existence theorem for strictly linear equations in $C^{2,\alpha}$ domains. This is an application of the Continuity Method (Theorem 5.3) and the proof relies on

- Kellogg's Theorem.
- Schauder's estimates.

Theorem 5.7 ([3], T6.8). *Let $\Omega \subset \mathbb{R}^n$ be a $C^{2,\alpha}$ domain and L a strictly elliptic linear operator in Ω with coefficients in $C^\alpha(\bar{\Omega})$, with $c \leq 0$. Assume that the Dirichlet problem for Poisson's equation $\Delta u = f$ in Ω , $u = \varphi$ on $\partial\Omega$ has a $C^{2,\alpha}(\bar{\Omega})$ solution for all $f \in C^\alpha(\bar{\Omega})$ and all $\varphi \in C^{2,\alpha}(\bar{\Omega})$. Then the Dirichlet problem*

$$(101) \quad \begin{aligned} Lu &= f \text{ in } \Omega \\ u &= \varphi \text{ on } \partial\Omega, \end{aligned}$$

also has a unique solution in $C^{2,\alpha}(\bar{\Omega})$ for all such f and φ .

Proof. Note that (by linearity) we can restrict to the case of zero boundary values: $Lv = g$ in Ω and $v = 0$ on $\partial\Omega$ (take $v = u - \varphi$, and $g = f - L\varphi$).

Now the following family of equations is considered

$$(102) \quad L_t = (1 - t)\Delta + tL, \quad 0 \leq t \leq 1,$$

where from the hypothesis on the coefficients of L we can see that L_t satisfies

$$(103) \quad \begin{aligned} a_t^{ij}(x)\xi_i\xi_j &\geq \lambda_t|\xi|^2, \quad \forall x \in \Omega, \xi \in \mathbb{R}^n, \\ |a_t^{ij}|_{C^\alpha(\Omega)}, |b_t^i|_{C^\alpha(\Omega)}, |c_t|_{C^\alpha(\Omega)} &\leq \Lambda_t. \end{aligned}$$

These bounds are independent of t when we take

$$(104) \quad \lambda_t = \min(1, \lambda), \quad \Lambda_t = \max(1, \Lambda).$$

Let $B_1 = \{u \in C^{2,\alpha}(\Omega) : u = 0, \text{ on } \partial\Omega\}$ and $B_2 = C^\alpha(\bar{\Omega})$. Note then that $L_t : B_1 \rightarrow B_2$ is a bounded linear operator between Banach spaces. Let u_t a solution of $L_t u = f$ for arbitrary $f \in C^\alpha(\Omega)$. From the maximum principle, we obtained Theorem 5.1, and in this case we have

$$(105) \quad |u_t|_{C^0(\Omega)} \leq C|f|_{C^\alpha(\Omega)},$$

From (105) and the Schauder's estimates (74) in Theorem 5.2 we have

$$(106) \quad |u_t|_{C^{2,\alpha}(\Omega)} \leq C(|u_t|_{C^0(\Omega)} + |f|_{C^\alpha(\Omega)}) = C|f|_{C^\alpha(\Omega)},$$

equivalently we have

$$(107) \quad |u_t|_{B_1} \leq C|L_t u|_{B_2}$$

with C is independent of t . If the domain is the ball $\Omega = B$, then L_0 is onto will follow from Kellogg's theorem (Theorem 4.13), which states that the solution u of the Dirichlet problem for the Poisson's equation belongs to $C^{2,\alpha}(\bar{B})$ when $f \in C^\alpha(\bar{B})$ and $\varphi \in C^{2,\alpha}(\bar{B})$. In our case, this is part of our hypothesis, hence we have that $L_0 = \Delta : B_1 \rightarrow B_2$ is onto and by the continuity method L_1 is invertible. \square

It is possible to develop the regularity theory in order to drop the assumption that Kellogg's Theorem hold for more general domains Ω .

Theorem 5.8 ([3], T6.14). *Let L be strictly elliptic in a bounded domain Ω with $c \leq 0$ and let f and the coefficients of L belong to $C^\alpha(\bar{\Omega})$. Suppose that Ω is a $C^{2,\alpha}$ domain and that $\varphi \in C^{2,\alpha}(\bar{\Omega})$. If $u \in C^0(\bar{\Omega}) \cap C^{2,\alpha}(\Omega)$ is a solution of the Dirichlet problem $Lu = f$ in Ω , $u = \varphi$ on $\partial\Omega$. Then $u \in C^{2,\alpha}(\bar{\Omega})$.*

Proof. Since Ω is in particular a C^2 domain we apply Theorem 5.6. There exist then a solution $u \in C^0(\bar{\Omega}) \cap C^{2,\alpha}(\Omega)$. Then we have to check only that at any point $x_0 \in \partial\Omega$ there is a neighbourhood V of x_0 such that $u \in C^{2,\alpha}(V \cap \bar{\Omega})$. \square

To investigate higher regularity of solutions, the following lemma states that if u is a $C^2(\Omega)$ solution of $Lu = f$, and if f and the coefficients of L are in C^α , then u should also be in $C^{2,\alpha}$.

Lemma 5.9 ([3], L6.16). *Let $u \in C^2(\Omega)$ be a solution of the equation $Lu = f$, in an open set Ω , where f and the coefficients of the elliptic operator L are in $C^\alpha(\Omega)$. Then $u \in C^{2,\alpha}(\Omega)$.*

The previous lemma and Schauder estimates gives the following

Theorem 5.10 (Interior Regularity Theorem ([3], T6.17)). *Let $u \in C^2(\Omega)$ be a solution of the equation $Lu = f$ in the open set Ω , where f and the coefficients of the elliptic operator L are in $C^{j,\alpha}(\Omega)$. Then $u \in C^{2+j,\alpha}(\Omega)$.*

Proof. For $j = 0$, we have the previous lemma. Let $j = 1$, so f and the coefficients of L are in $C^{1,\alpha}(\Omega)$, and assume $u \in C^2(\Omega)$. Hence $u \in C^{2,\alpha}(\Omega)$. Define for $h \in \mathbb{R}$ small, and $e_1 = (1, 0, \dots, 0) \in \mathbb{R}^n$ the following quotient

$$u^h := h^{-1}(u(x + he_1) - u(x)).$$

Note that

$$(108) \quad Lu^h = a_{ij}u_{ij}^h + b_i u_i^h + cu^h = f^h - a_{ij}^h u_{ij}(x + he_1) - b_i^h u_i(x + he_1) - c^h u(x + he_1).$$

The fact that $f \in C^{1,\alpha}(\Omega)$ and the following integral identity

$$f^h(x) = \frac{1}{h} \int_0^1 \frac{d}{dt} f(x + the_1) dt = \int_0^1 \frac{\partial}{\partial x_1} f(x + the_1) dt,$$

imply that $f^h \in C^\alpha(\Omega)$. Same conclusion applies for the coefficients a_{ij}^h, b_i^h, c^h . Considering $u^h \in C^2(\Omega)$ a solution to (108), the Schauder estimates implies that $u^h \in C^{2,\alpha}$ and using convergence arguments one shows that so is $\lim_{h \rightarrow 0} u^h(x) = \frac{\partial u(x)}{\partial x_1}$, and thus, that u is actually $C^{3,\alpha}$. The rest of the proof follows by induction. \square

Moreover, it is possible to obtain also the global regularity theorem using a stronger version of the lemma above and have a priori estimates up to the boundary

Theorem 5.11 (Global Regularity Theorem ([3], T6.19)). *Let Ω be a $C^{j,\alpha}$ domain, $j \geq 0$, and $\varphi \in C^{k+2,\alpha}(\bar{\Omega})$. Let $u \in C^0(\bar{\Omega}) \cap C^2(\Omega)$ be a solution of the Dirichlet problem $Lu = f$ in Ω and $u = \varphi$ on $\partial\Omega$, where f and the coefficients of the strictly (uniformly) elliptic operator L are in $C^{j,\alpha}(\bar{\Omega})$. Then $u \in C^{2+j,\alpha}(\bar{\Omega})$.*

6. QUASI-LINEAR SECOND ORDER PARTIAL DIFFERENTIAL EQUATIONS:
THE LERAY-SCHAUDER FIXED POINT THEOREM.

Following the same spirit of the last section, here we consider the Dirichlet problem $Qu = 0$ in Ω , $u = \varphi$ on $\partial\Omega$ such that Ω is a bounded domain, the coefficients of Q are C^α in their domain, the values on the boundary $\varphi \in C^{2,\alpha}(\bar{\Omega})$. Then the existence of a solution $u \in C^{2,\alpha}(\bar{\Omega})$ if we can establish the a priori estimate

$$(109) \quad |u|_{C^{1,\beta}(\bar{\Omega})} \leq M.$$

A four-step process is stated in [3] to get this estimate is the successive estimation of $\sup_{\Omega} |u|$ (using maximum principle for quasilinear equations), $\sup_{\partial\Omega} |Du|$, $\sup_{\Omega} |Du|$, each using the preceding ones, and finally obtain an estimate for $|u|_{C^{1,\beta}(\bar{\Omega})}$ for some $\beta > 0$.

There are lots of results regarding the gradient estimates in the interior and in the boundary. It is possible to obtain interior estimates in terms of the gradient estimate on the boundary by a refinement of the so called Bernstein technique.

The existence result is obtained by an extension of the Brouwer fixed point theorem:

Theorem 6.1. *Let $B \subset \mathbb{R}^n$ be an open ball and $T : \bar{B} \rightarrow \bar{B}$ a continuous map. Then there is some $x \in \bar{B}$ such that $T(x) = x$, i.e., T has at least one fixed point.*

Theorem 6.2 ([3], T11.3, Leray-Schauder). *Let B a Banach space and $T : B \rightarrow B$ a compact mapping, that is, the image under T of any bounded set have compact closure. Suppose that there exists $M > 0$ such that*

$$(110) \quad |x|_B < M$$

for all $x \in B$. Then for all $\sigma \in [0, 1]$, σT has a fixed point.

The following result is basic to the theory of second order quasilinear equations. Indeed, its discovery by De Giorgi [DG I] and Nash [NA] for operators of the form $Lu = D_i(a^{ij}(x)D_j u)$ essentially opened up the theory of quasilinear equations in more than two variables.

Theorem 6.3 ([3], T8.22). *Let the operator L be of the form*

$$(111) \quad Lu = D_i(a^{ij}(x)D_j u + b^i(x)u) + c^i(x)D_i u + d(x)u,$$

and uniformly elliptic with $0 < \lambda < \Lambda$. Let $f^i, g, i = 1, 2, \dots, n$ be locally integrable functions in Ω and consider the problem

$$(112) \quad Lu = g + D_i f^i, \quad \text{in } \Omega.$$

Assume in addition that $f^i \in L^q(\Omega)$ for $i = 1, 2, \dots, n$, and for some $q > n$, $g \in L^{q/2}(\Omega)$. If $u \in W^{1,2}(\Omega)$ is a solution to the equation in Ω , then u is Hölder continuous in Ω , and for any ball $B_0 = B_{R_0}(y) \subset \Omega$ and $R \leq R_0$ we have

$$(113) \quad \text{osc}_{B_R(y)} \leq CR^\alpha \left(R_0^{-\alpha} \sup_{B_0} |u| + k \right),$$

where the constant $C = C(n, \Lambda/\lambda, v, q, R_0)$ and $\alpha = \alpha(n, \Lambda/\lambda, v, R_0, q)$ are positive constants and $k = \lambda^{-1}(\|f\|_q + \|g\|_{q/2})$.

Now we apply this result to establish the existence of a solution for the quasi-linear case.

Theorem 6.4 ([3], T11.4). *Let $\Omega \in \mathbb{R}^n$ be a bounded domain and suppose that the quasilinear equation Q is elliptic in $\bar{\Omega}$ with Hölder continuous coefficients $a^{ij}, b \in C^\alpha(\bar{\Omega} \times \mathbb{R} \times \mathbb{R}^n)$, $0 \leq \alpha \leq 1$, the boundary $\partial\Omega \in C^{2,\alpha}$ and assume that the boundary values $\varphi \in C^{2,\alpha}(\bar{\Omega})$. Consider the following one parameter family of boundary value problems*

$$(114) \quad \begin{cases} Q_\sigma u = 0 & \text{in } \Omega \\ u = \sigma\varphi & \text{on } \partial\Omega, \end{cases}$$

where Q_σ is defined for $0 \leq \sigma \leq 1$ by

$$(115) \quad Q_\sigma u = a^{ij}(x, u, Du)D_{ij}u + \sigma b(x, u, Du).$$

If for some $\beta > 0$ there is a constant M independent of u and σ , such that for every solution $u \in C^{2,\alpha}(\bar{\Omega})$ of the Dirichlet problem $Q_\sigma u = 0$ in Ω , $u = \sigma\varphi$ on $\partial\Omega$, and this constant satisfies

$$(116) \quad |u|_{C^{1,\beta}(\bar{\Omega})} < M,$$

then the Dirichlet problem $Qu = 0$ in Ω , $u = \varphi$ on $\partial\Omega$, is solvable in $C^{2,\alpha}(\bar{\Omega})$.

Proof. Recall that if $0 < \alpha \leq \beta < 1$, and Ω is a bounded set, then for every $f \in C^\beta(\Omega)$ we have

$$(117) \quad |f|_{C^\alpha(\Omega)} \leq \text{diam } \Omega^{\beta-\alpha} |f|_{C^\beta(\Omega)},$$

from where the inclusion $C^\beta(\Omega) \rightarrow C^\alpha(\Omega)$ follows. This inclusion is also compact as consequence of Arzela-Ascoli theorem. Then there is a compact inclusion of $C^{2,\alpha\beta}(\bar{\Omega})$ in $C^{1,\beta}(\bar{\Omega})$ and another in $C^{2,\alpha}(\bar{\Omega})$.

Define the operator $T : C^{1,\alpha}(\bar{\Omega}) \rightarrow C^{2,\alpha\beta}(\bar{\Omega})$ by letting $Tv = u$ be the unique solution in $C^{2,\alpha\beta}(\bar{\Omega})$ of the linear Dirichlet problem,

$$(118) \quad \begin{aligned} Qu &= a^{ij}(x, v, Dv) D_{ij}u + b(x, v, Dv) = 0, \text{ in } \Omega \\ u &= \varphi, \text{ on } \partial\Omega, \end{aligned}$$

where we are applying the existence and uniqueness result from the theory of elliptic linear equations. Then a solution in $u \in C^{2,\alpha}(\bar{\Omega})$, is a fixed point of T , for a solution of the Dirichlet problem above is actually a solution of the equation $Tu = u$ in $C^{1,\beta}(\bar{\Omega})$.

Note on the other hand that the equation $\sigma Tu = u$ in $C^{1,\beta}(\bar{\Omega})$, is equivalent to the Dirichlet problem

$$(119) \quad \begin{aligned} Q_\sigma u &= a^{ij}(x, u, Du) D_{ij}u + \sigma b(x, u, Du) = 0, \text{ in } \Omega \\ u &= \sigma\varphi, \text{ on } \partial\Omega, \end{aligned}$$

We have to show the compactness and continuity of the operator T in order to apply the Leray-Schauder theorem. That T is a compact mapping follows from the fact that any bounded set in $A \subset C^{1,\beta}(\bar{\Omega})$, then by Schauder's estimate, T maps A into a bounded set in $C^{2,\alpha\beta}(\bar{\Omega})$. By Arzela's Theorem, this is precompact in $C^2(\bar{\Omega})$ and $C^{1,\beta}(\bar{\Omega})$.

Continuity of T we take a convergent sequence $\lim v_m = v$ in $C^{1,\beta}(\bar{\Omega})$. Note that Tv_m is precompact in $C^2(\bar{\Omega})$. This means that for every subsequence of $\{v_m\}$, the corresponding $\{Tv_m\}$ has convergent subsequence. Call $\{T\bar{v}_m\}$ such a convergent subsequence and let $u \in C^2(\bar{\Omega})$ be its limit. Since

$$(120) \quad \begin{aligned} 0 &= \lim_{n \rightarrow \infty} \{a^{ij}(x, \bar{v}_m, D\bar{v}_m) D_{ij}T\bar{v}_m + b(x, \bar{v}_m, D\bar{v}_m)\} \\ &= a^{ij}(x, v, Dv) D_{ij}u + b(x, v, Dv), \end{aligned}$$

then we conclude that $Tv = u$ and then the sequence Tv_m converges to u . \square

7. FULLY NONLINEAR PARTIAL DIFFERENTIAL EQUATION

We are concern in operators of the type

$$(121) \quad F[u] = F(x, u, Du, D^2u)$$

where $\Omega \subset \mathbb{R}^n$ is an open domain, $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$, Du denote the gradient of u and D^2u the hessian matrix. F is then a function on $\Gamma = \Omega \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^{n \times n}$, where $\mathbb{R}^{n \times n}$ denotes the space of real symmetric $n \times n$ matrices and write $\gamma = (x, z, p, r) \in \Gamma$. If the function F is an affine function with respect to the r variables then we say that F is *quasilinear*, and in any other case we say that F is *fully nonlinear*.

Definition 7.1. The operator F is called **elliptic** in $\mathcal{U} \subset \Gamma$ if the matrix

$$(122) \quad F_{ij}(\gamma) = \frac{\partial F}{\partial r_{ij}}(\gamma)$$

is positive. If $\lambda(x)$ and $\Lambda(x)$ denote respectively the minimum and maximum eigenvalue of $[F_{ij}(x)]$, then F is called **uniformly elliptic** if Λ/λ is bounded and **strictly elliptic** if $1/\lambda$ is bounded.

Example 7.2. One can see that in dimension $n = 2$, the ellipticity of $F = F(x, u, Du, D^2u)$ is equivalent to have

$$(123) \quad 4F_{11}F_{22} - F_{12}^2 > 0.$$

One example to have in mind is the Monge-Ampère equation

$$(124) \quad \det(D^2u) = f \text{ in } \Omega.$$

It turns out to be elliptic in the class of convex functions u , and then necessarily $f > 0$.

We will see in this section that by the continuity method, the existence of a solution for the Dirichlet problem is reduced to obtain the a priori estimate

$$(125) \quad |u|_{C^{2,\alpha}(\bar{\Omega})} \leq C,$$

for some $0 < \alpha < 1$. Then, like in the quasilinear case we have to establish estimates for $\sup_\Omega |u|$, $\sup_{\partial\Omega} |Du|$, $\sup_\Omega |Du|$, and additionally $\sup_{\partial\Omega} |D^2u|$, $\sup_\Omega |D^2u|$.

Int he case F is an uniformly elliptic, fully nonlinear concave equation, the Evans-Krylov theorem [1][4], gives the following a priori estimate

$$(126) \quad |u|_{C^{2,\alpha}(\bar{\Omega})} \leq C |u|_{C^{1,1}(\bar{\Omega})},$$

when for the case $\Omega = B_1$ is the unit ball and C depends only on the concavity of F .

8. MAXIMUM PRINCIPLE FOR NONLINEAR PDE'S

Recall that in the case of the Laplacian, two solutions u_1 and u_2 of the equation $\Delta u = 0$ in Ω . Then if the difference $u_1 - u_2$ is positive at some interior point of Ω , then it cannot become zero at some interior point of Ω .

One way to understand this fact is by noticing that the function

$$(127) \quad F(D^2u) = \Delta u = \text{Trace}[D^2u],$$

is a monotone function of the Hessian matrix $\left(\frac{\partial^2 u}{\partial x_i \partial x_j}\right)$. Then a natural family of equations to be consider are

$$(128) \quad F(D^2u) = 0,$$

where F is a strictly monotone function of D^2u .

In differential geometry, equations of these types appear, for instance, the coefficients of the characteristic polynomial of the Hessian.

$$(129) \quad P(\lambda) = \det(D^2u - \lambda I)$$

are such equations where D^2u is restricted to stay in the appropriate set of symmetric matrices $\mathbb{R}^{n \times m}$.

Then the following are examples that we should have in mind: If λ_i denotes the eigenvalues of D^2u then

- (1) Laplace: $\sigma_1 = \Delta u = \sum_i \lambda_i$.
- (2) $\sigma_2 = \sum_{i \neq j} \lambda_i \lambda_j$.
- (3) Monge-Ampere: $\sigma_n = \det(D^2u) = \prod_i \lambda_i$.

Note however that for the Monge-Ampere equation, the determinant is a monotone function of the Hessian provided that all eigenvalues λ_i 's are positive, i.e., provided that the functions under considerations are convex.

In general the setting is the following: We are interested in finding a function $u(x)$ in a domain $\Omega \subset \mathbb{R}^n$ such that

$$(130) \quad F(D^2u) = f(x), \quad \text{in } \Omega,$$

and with prescribed boundary data

$$(131) \quad u(x) = g(x), \quad \text{on } \partial\Omega,$$

And we have to assume that the function $F(M)$ is a function in the space of symmetric matrices such that

- (1) F is Lipschitz
- (2) F is strictly increasing in the direction of *positive matrices*.

If you put together these two conditions on F , then we can say that if $N > 0$ then

$$(132) \quad F(M) + c_1 \|N\| \leq F(M + N) \leq F(M) + c_2 \|N\|.$$

If we have $F(M + N) = F(M)$ for $N = N^+ + N^-$, then the positive part and the negative part should be comparable

$$(133) \quad \|N^+\| \sim \|N^-\|.$$

In the case of $F(\lambda_1, \lambda_2) = C$ then the level sets of F can be drawn

The main ingredients of the regularity theory for these class of fully nonlinear elliptic equations are

- The Aleksandrov-Bakelman-Pucci Theorem.
- Krylov-Safonov Harnack inequality for concave F .
- Evans-Krylov $C^{2,\alpha}$ - Theorem.

The structure of the theory

If we have translation invariant operators (like constants) then the first derivatives usually satisfy a nice equation, for instance the Linearised equation: Let $D_v u = Du \cdot v$ the derivative in the direction v , then

$$(134) \quad 0 = D_v F(D^2u(x)) = F^{ij}(D^2u) D_{ij}(D_v u).$$

Then $D_v u$ satisfies this equation, for which at least we know that the matrix F^{ij} is strictly positive definite and bounded, hence we can think of an equation of the form

$$(135) \quad 0 = a^{ij}(x) D_{ij}(w),$$

and conclude by Krylov-Safonov that w is Hölder and that they satisfy a Harnack inequality. This is the same strategy as when De Giorgi theorem and for equations in divergence form. This is where one can get if concavity is not assumed.

In the case F is also concave, then we take second order derivative of the equation and we get

$$(136) \quad 0 = D_{vv} F(D^2u(x)) = F^{ijkl}(D^2u) D_{ij}(D_v u) D_{kl}(D_v u) + F^{ij}(D^2u) D_{ij}(D_{vv} u),$$

and the concavity implies that

$$(137) \quad F^{ij,kl}(D^2u)D_{ij}(D_vu)D_{kl}(D_vu) \leq 0.$$

Then second derivatives of u are subsolutions of some equation, that is

$$(138) \quad F^{ij}(D^2u)D_{ij}(D_{vv}u) = a^{ij}(x)D_{ij}w \geq 0,$$

hence w is a subsolution that is bounded above (half of Harnack) then u is called *semi-concave* in the sense that

$$(139) \quad u - C|x|^2,$$

is concave.

Intermedian steps in Evans-Krylov: u is $C^{1,1}$ which implies that $D_{ij}u$ is bounded. The Evans-Krylov theorem Take a sub solution of an equation which is bounded (say by 1) And there is a “chunk” in the domain where it goes down to something strictly below. Then that pulls the subsolution all down. In other words, you pass from some information in measure, then you go to a ball of radius one half and what was an information in measure becomes an information in L infinity norm.

You start in the ball $B_1(x_0)$ and look at the its image D^2u . $T : B_1(x_0) \rightarrow \mathbb{R}^{n \times n}$; $x \mapsto D^2u(x)$. This lives in one of the level sets of F . Cover this level set (surface) with balls and then take them back to see where do they come from. Notice that one of the balls should come from a non-trivial set of positive measure where you can estimate what should be its measure of the best (largest) set.

If one of these ball has diameter smaller than the diameter of the image, then ther is some directional derivative that goes in the growing direction, then there is D^2u that at this h is bigger than here. But D^2u is a subsolution bounded above by the line, then ther is a chunk if you come back here that is not trivial, then the property of subsolution says that if you go to the ball of radius one half then the bound has to come down, this means that you have to reduce the diameter of the image, iteration scheme gives you that the diameter of image is C^α of the second derivative.

In the Monge-Ampere equaiton case and the σ_k equation, we want to find a geometric configuration where the theory applies. Consider the Monge-Ampere equation equal to a constant. In the first quadrant ($\lambda_j \geq 0$). Their product is monotone as long as all λ_i 's are strictly positive. Then, at the boundary of the first quadrant degenerates. So we want to stay inside the first quadrant. Consider

$$(140) \quad \det(D^2u) = 1.$$

The the level surface

$$(141) \quad \det(M) = 1$$

becomes vertical as $M \rightarrow \infty$. Note that if $\prod_i \lambda_i = 1$ and $\|\lambda\|$ is larfe, then must be λ_1 small and λ_2 large. Then $\prod_i \lambda_i = 1$ is small and a vertical perturbation in the λ_2 direction changes

$$(142) \quad \prod_i \lambda_i$$

only a little. Then we must have λ bounded to remain strictly elliptic.

From the concavity of $(\det(D^2u))^{1/2}$, second derivatives are subsolutions. If we can control them at the boundary, we are done since we are assumming Ω strictlyconvex and smooth, boundary data and right hand side smooth).

Theorem 8.1. *If $F(D^2u, x)$ is uniformly elliptic, i.e., if F is strictly monotone as a function of the Hessian, or in differential form, if*

$$(143) \quad F^{ij}(M) = \frac{\partial F}{\partial m_{ij}},$$

is uniformly positive definite, then the solutions of $F(D^2u) = 0$ are $C^{1,\alpha}$.

8.1. The classical Aleksandrov-Bakelman-Pucci Theorem.

Theorem 8.2. *Let u be a viscosity super solution of the linear equation*

$$(144) \quad Lu := a^{ij}(x)u_{ij}(x).$$

with

$$(145) \quad \begin{aligned} Lu &\leq f(x), & x \in B_1, \\ u(x) &\geq 0, & x \in \partial B_1. \end{aligned}$$

Suppose that the coefficients of $a^{ij}(x)$ are measurable functions and that the equation is uniformly elliptic, that is, there are positive constants $0 < \lambda < \Lambda$ such that for all $\xi \in \mathbb{R}^n$ it holds

$$(146) \quad \lambda|\xi|^2 \leq a^{ij}(x)\xi_i\xi_j \leq \Lambda|\xi|^2.$$

Assume also that the function f is continuous. Then the ABP theorem states that

$$(147) \quad \sup_{B_1} |(u^-)|^n \leq C \int_{u=\Gamma_u} (f^+)^n dx,$$

where $C = C(n, \lambda, \Lambda)$, and Γ_u is the convex envelope of u , that is, it is the largest non-positive convex function in B_2 that lies above u in B_1

An important fact to notice is that the integration on the right hand side takes place only on the set where u agrees with its convex envelope is an important feature of the estimate and it is not to be overlooked.

8.2. Krylov-Safonov theorem.

Theorem 8.3. Suppose that u is a bounded solution of the following uniformly elliptic PDE

$$(148) \quad a^{ij}(x)u_{ij}(x) + b^i u_i(x) = f(x) \quad x \in B_1,$$

with ellipticity constants $0 < \lambda < \Lambda$, and coefficients $b^i, f \in L^n(B_1)$. Then the function u is Hölder continuous, and for some $0 < \alpha < 1$ it holds

$$(149) \quad |u|_{C^\alpha(B_{1/2})} \leq C(|u|_{L^\infty(B_1)} + |f|_{L^n(B_1)}),$$

where $C = C(n, \lambda, \Lambda, |b|_{L^\infty})$.

Theorem 8.4. Suppose that u is a bounded non-negative solution of the following uniformly elliptic PDE

$$(150) \quad a^{ij}(x)u_{ij}(x) + b^i u_i(x) = f(x) \quad x \in B_1,$$

with ellipticity constants $0 < \lambda < \Lambda$, and coefficients $b^i, f \in L^n(B_1)$. Then the following Harnack inequality holds:

$$(151) \quad \sup_{B_{1/2}} u \leq C \left(\inf_{B_{1/2}} u + |f|_{L^n(B_1)} \right),$$

where $C = C(n, \lambda, \Lambda, |b|_{L^\infty})$.

Theorem 8.5. Suppose that u is a bounded viscosity solution of the fully nonlinear elliptic equation

$$(152) \quad F(D^2 u) = 0, \quad \text{in } B_1.$$

Then there exists $0 < \alpha < 1$, $\alpha = \alpha(n, \lambda, \Lambda)$, such that $u \in C^{1,\alpha}(B_1)$, and

$$(153) \quad |u|_{C^{1,\alpha}(B_1)} \leq C(|u|_{L^\infty(B_1)} + F(0)).$$

The constant $C = C(n, \lambda, \Lambda)$.

8.3. Evans - Krylov.

Theorem 8.6 ([3], T17.14). Let $u \in C^4(\Omega)$ with $F[u] = 0$ in Ω . Assume that $F \in C^2(\Gamma)$ is uniformly elliptic, F is a concave function on $r = D^2 u$. Then for any $\Omega' \subset \subset \Omega$ it holds

$$(154) \quad \|D^2 u\|_{C^\alpha(\Omega')} \leq C,$$

where α depends on n, λ, Λ and C depends in addition on $\|u\|_{C^2(\Omega)}$, $\text{dist}(\Omega', \Omega)$ and the first and second derivatives of F other than F_{rr} .

Theorem 8.7 ([3], T17.26'). Let $\Omega \subset \mathbb{R}^n$ bounded with boundary $\partial\Omega \in C^3$, and let $\varphi \in C^3(\overline{\Omega})$. Suppose that $u \in C^3(\overline{\Omega}) \cap C^4(\Omega)$ is a solution of $F[u] = 0$ in Ω , $u = \varphi$ on $\partial\Omega$. Assume that $F \in C^2(\overline{\Gamma})$ is uniformly elliptic, F is a concave function on $r = D^2 u$. Then we have the following estimate

$$(155) \quad \|u\|_{C^{2,\alpha}(\Omega)} \leq C,$$

where α depends on n, λ, Λ and C depends in addition on $\|u\|_{C^2(\Omega)}$, $\text{dist}(\Omega', \Omega)$ and the first and second derivatives of F other than F_{rr} .

REFERENCES

- [1] L. C. Evans. Classical solutions of fully nonlinear, convex, second-order elliptic equations. *Comm. Pure Appl. Math.*, 35(3):333–363, 1982.
- [2] L. C. Evans. *Partial differential equations*. American Mathematical Society, 2010.
- [3] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [4] N. V. Krylov. Boundedly inhomogeneous elliptic and parabolic equations. *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(3):487–523, 670, 1982.
- [5] L. Nirenberg. *Lectures on Differential Equations and Differential Geometry*. Classical Topics in Mathematics 7. Higher Education Press, 2019.